

Theoretical Study of Friction: A Case of One-Dimensional Clean Surfaces.

Hiroshi Matsukawa and Hidetoshi Fukuyama+

Department of Physics, Faculty of Science, Nara Women's University,
Nara 630, Japan

Department of Physics, Faculty of Science, Osaka University,

1-16, Machikaneyama-cho, Toyonaka, Osaka 560, Japan* +Department of Physics, Faculty of Science,
University of Tokyo,

7-3-1, Hongo, Bunkyo-ku, Tokyo 113, Japan

(Received 30 November 1993) Abstract A new method has been proposed to evaluate the frictional force in the stationary state. This method is applied to the 1-dimensional model of clean surfaces. The kinetic frictional force is seen to depend on velocity in general, but the dependence becomes weaker as the maximum static frictional force increases and in the limiting case the kinetic friction gets only weakly dependent on velocity as described by one of the laws of friction. It is also shown that there is a phase transition between state with vanishing maximum static frictional force and that with finite one. The role of randomness at the interface and the relation to the impurity pinning of the sliding Charge-Density-Wave are discussed.

PACS numbers: 46.30.P, 81.40.P, 68.35.J

*Present Address e-mail address: hiro@ena.wani.osaka-u.ac.jp To appear in Phys.Rev.B.

1. Introduction

The study of friction has been carried out for a long time¹. Recently the developments of new technologies enable us to investigate the friction in a situation which was not considered before²⁻¹⁰. It is well known that the three laws of friction hold well in a usual situation¹: (i) The frictional force does not depend on an apparent area of contact surfaces. (ii) The frictional force is proportional to the normal load, where the proportionality constant is called the coefficient of friction. (iii) The kinetic frictional force does not depend on the relative velocity of contact surfaces and is less than the maximum static frictional force. It is also known that there are cases where these laws of friction no longer hold. The limitation of the laws and the possible new laws which hold beyond such limitation have, however, not been clarified to the best of our knowledge. Actually even though there exists attempts of the theoretical explanation on the laws of static friction as discussed below, the concrete explanation of the laws of kinetic friction is lacking. This paper is our first step to study these basic questions on friction theoretically.

Theoretical explanation of the laws of static friction have been as follows¹. Due to the surface roughness the area of actual contacting points is much less than the apparent contact area and the pressure there reaches the yield pressure. Then the total area of actual contact points is proportional to the normal load. The two surfaces adhere to each other by intermolecular forces at the contact. The maximum static frictional force is equal to the shear strength of the contact times the total area of contacting points and is then proportional to the normal load. This view can not be applied to the clean surface without surface randomness⁶. For example, as the interatomic force between two bodies works among all atoms at surfaces, the frictional force is expected to survive even for the vanishing normal load and to be proportional to the contacting area and then the first and second laws of friction will no longer hold.

Recently Hirano and Shinjo⁶⁻⁸ pointed out another peculiar feature of the friction of clean surfaces. That is the frictional transition, which is the phase transition between states with and without finite maximum static frictional force. In the simplest case of 1-dimension(d) where the atoms of the one body are fixed, the model reduces to the Frenkel-Kontrova model¹¹. In that model such a phase transition is known to exist when the ratio of mean atomic distances of two bodies is irrational, i.e., in the incommensurate case. This transition is called Aubry's breaking of analyticity transition¹². When the strength of interatomic potential is less than a critical value the spatial configuration of atoms in the ground state is smooth and the maximum static frictional force vanishes. Above the critical value the configuration has discommensurate structure and the maximum static frictional force is finite. Hirano and Shinjo have claimed the existence of such a frictional transition even in a 3d model where the atoms of the lower body are fixed, and that the static frictional force can vanish between pure metals with incommensurate clean surfaces, e.g., (111) and (110) surfaces of α -iron^{6,8}.

All of these studies are, however, based on the model where atoms in one of the interfaces have held fixed. Nothing is known about the frictional transition in a more realistic model where both atoms can relax. The behavior of kinetic friction has also not been explored yet.

Deferring the detailed studies on rough surfaces until future publications, we study in this paper the static and kinetic frictional forces of the 1d model of clean surfaces, where atoms in both sides of the interface can relax. In the theoretical study of friction, it is not clear even what should be calculated as the frictional force and a method of calculation has not been established. We first propose a new method to calculate the kinetic frictional force in a stationary state and then apply it to 1d model of clean surfaces. We derive the explicit velocity dependence of the kinetic friction of the present microscopic model. It is seen that the velocity dependence is appreciable in general. This dependence becomes weaker as the maximum static frictional force increases and eventually the kinetic friction gets almost velocity independent as described by the third law of friction. At the same time we also find the frictional transition in the present model.

This paper is organized as follows. In 2 we address ourselves to the method to calculate the frictional force. 3 we define the 1d model of clean surfaces and discuss the relationship of the present model with the Frenkel-Kontrova model. Then the numerical results based on the model are presented. Finally in .4 we summarize the present results and discuss the effects of randomness and the relation to the impurity pinning of the sliding Charge-Density-Wave.